

USING COLORS TO IMPROVE PHOTOMETRIC METALLICITY ESTIMATES FOR GALAXIES

N. E. SANDERS¹, E. M. LEVESQUE^{2,3}, A. M. SODERBERG¹

Draft version October 23, 2012

ABSTRACT

There is a well known correlation between the mass and metallicity of star-forming galaxies. Because mass is correlated with luminosity, this relation is often exploited, when spectroscopy is not available, to estimate galaxy metallicities based on single band photometry. However, we show that galaxy color is typically more effective than luminosity as a predictor of metallicity. This is a consequence of the correlation between color and the galaxy mass-to-light ratio and the recently discovered correlation between star formation rate (SFR) and residuals from the mass-metallicity relation. Using Sloan Digital Sky Survey spectroscopy of 148,021 nearby galaxies, we derive “*LZC* relations,” empirical relations between metallicity (in nine common strong line diagnostics), luminosity, and color (in three filter pairs). We show that these relations allow photometric metallicity estimates, based on luminosity and a single optical color, that are $\sim 40\%$ more precise than those made based on luminosity alone; galaxy metallicity can be estimated to within $\sim 0.06 - 0.1$ dex of the spectroscopically-derived value depending on the metallicity diagnostic used. Including color information in metallicity estimates also reduces systematic biases for populations skewed toward high or low SFR environments, as we illustrate using the host galaxy of the supernova SN 2010ay. This new tool will lend more statistical power to studies of galaxy populations, such as supernova and gamma-ray burst host environments, in ongoing and future wide field imaging surveys.

Subject headings: galaxies: abundances — galaxies: photometry — galaxies: ISM

1. INTRODUCTION

The gas-phase metallicity of galaxies, as measured from their nebular emission spectrum, is correlated with galaxy luminosity (Lequeux et al. 1979; Garnett & Shields 1987). This relation has been used as a key observational tool in the study of populations such as supernova host galaxies (e.g. Prantzos & Boissier 2003; Arcavi et al. 2010), where gas-phase metallicity is an important proxy for the properties of their short lived progenitor stars. However, using Sloan Digital Sky Survey (SDSS) imaging and spectroscopy for $\sim 53,000$ galaxies, Tremonti et al. (2004) showed that the luminosity-metallicity relation has a large intrinsic scatter of $\sigma = 0.16$ dex (50%), in terms of metallicity residuals, which limits the utility of this relation as an effective indicator of metallicity.

There are two primary causes for the scatter in the luminosity-metallicity relation. First, while the scatter in the mass-metallicity relation is fairly small ($\sigma = 0.10$ dex, Tremonti et al. 2004), luminosity is not a perfect proxy for mass. The mass-to-light ratio of galaxies is highly correlated with galaxy color, such that redder galaxies at a fixed luminosity are more massive (Bell & de Jong 2001; Kauffmann et al. 2003b). Second, a more fundamental relation has been uncovered between mass (M), metallicity (Z), and star formation rate (SFR) (Lara-López et al. 2010; Mannucci et al. 2010). This “fundamental plane” or “Fundamental Metallicity Relation” has remarkably small residual scatter ($\sigma = 0.05$ dex), indicating that

variations in SFR are responsible for much of the scatter in the mass-metallicity relation. The existence of this fundamental plane is a valuable constraint for models of galaxy evolution, and likely an expression of galaxy outflows, infall, and/or downsizing (Mannucci et al. 2010; Peeples & Shankar 2011). To improve the precision of photometric metallicity estimates, a readily accessible observable must be used to break the degeneracy between luminosity, mass, and SFR.

In this paper, we show that the addition of color information significantly decreases the scatter in photometric metallicity estimates. We derive the optimal projection of the fundamental plane for star-forming galaxies, in terms of the observable properties luminosity and color, that we call the *LZC* relation. We describe the sample of SDSS galaxies we use to study these correlations and methods for spectroscopic metallicity estimation in Section 2. In Section 3, we derive analytic expressions for the *LZC* as expressed in a variety of different filter sets and metallicity diagnostics. Finally, in Section 4, we describe how specific observational studies may benefit from the *LZC* relation in making precise metallicity determinations from imaging available from wide field sky surveys.

2. GALAXY SAMPLE

We used the MPA/JHU catalog⁴ of 927,552 star-forming galaxies from the SDSS-DR7 (Abazajian et al. 2009). The catalog includes synthesized rest-frame photometry corrected for Galactic extinction, emission line fluxes, stellar masses (based on SED fitting to *ugriz* photometry), and star formation rates for each galaxy, as described in Kauffmann et al. (2003b); Brinchmann et al.

nsanders@cfa.harvard.edu

¹ Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138 USA

² CASA, Department of Astrophysical and Planetary Sciences, University of Colorado, 389-UCB, Boulder, CO 80309, USA

³ Einstein Fellow

⁴ The MPA/JHU catalog is available at <http://www.mpa-garching.mpg.de/SDSS>

TABLE 1
METALLICITY DIAGNOSTICS USED

Name	Line Ratios	Source
M91	R23	McGaugh 1991 ^a
P01	P	Pilyugin 2001
KD02	N2O2	Kewley & Dopita 2002 ^b
KK04	R23	Kobulnicky & Kewley 2004
PP04	N2,O3N2	Pettini & Pagel 2004
T04	model ^c	Tremonti et al. 2004
PT05	P	Pilyugin & Thuan 2005
PVT	P	Pilyugin et al. 2010 ^d

^a We have adopted the revised prescription suggested by Kobulnicky et al. (1999).

^b Following Kewley & Ellison (2008), we adopt the average of the M91 and KK04 for the lower branch solution.

^c The T04 metallicities are estimated based on simultaneous fits of all major emission lines to photoionization models and are provided in the MPA/JHU catalog

^d We use the “ONS” solution, which includes a dependence on the S II flux, for the conditions $\log(\text{N2}) > -0.1$ and $\log(\text{N2}/\text{S2}) > -0.25$ (which is true for $\sim 98\%$ of the SDSS galaxies).

(2004); Salim et al. (2007). We correct line fluxes for line-of-sight extinction (including both Galactic and intrinsic reddening) from the Balmer flux decrement. We assume $F_{\text{H}\alpha}/F_{\text{H}\beta} = 2.85$ (corresponding to $T = 10,000$ K and $n_e = 10^4 \text{ cm}^{-3}$ for Case B recombination; Osterbrock & Ferland 2006) and the extinction curve of Cardelli et al. (1989), assuming $R_V = 3.1$.

We perform preliminary sample cuts on the catalog following the prescription of Mannucci et al. (2010), as follows. We require $0.07 < z < 0.3$, $(\text{S/N})_{\text{H}\alpha} > 25$, $F_{\text{H}\alpha}/F_{\text{H}\beta} > 2.5$, $A_V < 2.5$ mag, and we reject AGN following Kauffmann et al. (2003a). In total, following these cuts, we consider 148,021 galaxies.

For each galaxy, we compute oxygen abundance as a proxy for metallicity using a variety of strong line diagnostics that are widely used in the literature (Table 1; see Lopez-Sanchez et al. 2012 for a recent review). First, we employ several diagnostics relying on the R23 ratio, which depends on the fluxes of O II $\lambda\lambda 3726, 3729$, O III $\lambda 4959$ and $\lambda 5007$, and H β . The R23 diagnostic suffers from a degeneracy (see e.g. Kewley & Dopita 2002) that we break using either the N2O2 (N II $\lambda 6584$ to O II $\lambda 3727$) or N2 (N II $\lambda 6584$ to H α) ratios, following the authors’ prescriptions. Next we employ diagnostics depending on N2, N2O2, and O3N2, the flux ratio of O III $\lambda 5007$ and N II $\lambda 6584$. Finally, we employ diagnostics based on the ionization parameter, P , the ratio of O III $\lambda 4959$ and $\lambda 5007$ to R23. It is necessary to use multiple diagnostics because they exhibit well known systematic discrepancies, which are particularly strong between those diagnostics calibrated empirically and those calibrated against photoionization models (Kewley & Ellison 2008).

Figure 1 demonstrates two correlations in the galaxy sample. First, it shows the well known luminosity-metallicity correlation, where more luminous galaxies (shown in M_r) typically have higher metallicities. However, there is significant scatter in this relation, with a standard deviation of $\sigma_0 = 0.14$ dex in the residuals in metallicity for r band, as shown. Second, the figure demonstrates that there is a correlation between the




FIG. 1.— A luminosity-metallicity (r -band, T04) plot showing a random subset of 10,000 SDSS galaxies, color coded by $g-r$ color. The solid lines show the median luminosity-metallicity relation for the galaxies divided into 10 equal-sized bins in $g-r$ color.

residual in metallicity (the offset from the luminosity-metallicity relation) and galaxy color.

3. LUMINOSITY-METALLICITY-COLOR RELATION

Following Mannucci et al. (2010), we project the LZC relation onto an axis μ with components of color and luminosity:

$$\mu = M_i - \alpha \times (m_j - m_k) \quad (1)$$

$$12 + \log(\text{O}/\text{H}) = p_0 + p_1 \mu + p_2 \mu^2 + p_3 \mu^3 \quad (2)$$

where i, j, k are choices of filters and p_l are parameters of a third order polynomial. For each combination of metallicity diagnostic, luminosity band, and color, we determine the optimal projection of the galaxy LZC relation by sampling from a grid of α parameters and selecting the value that minimizes the variance in the residuals of metallicity. We report the optimal value of α and corresponding best fit LZC parameters p in Table 2.

We find a correlation between the luminosity-metallicity relation and color which varies in strength depending on the choice of metallicity diagnostic and filter set. Figure 2 demonstrates this optimization for one filter set and three choices of metallicity diagnostic: T04, KD02, and PP04 O3N2. For PP04 O3N2, the scatter in the metallicity residuals of the LZC relation is $\sigma_Z = 0.10$ dex ($\sim 25\%$), as compared to the LZ relation ($\alpha = 0$), $\sigma_{Z,0} = 0.13$ dex ($\sim 35\%$). The decrease in residual scatter is similar in other diagnostics, ranging from $\sigma_Z/\sigma_{Z,0} = 0.68 - 0.87$, with the exceptions of P01 and PT05 (discussed below). Note that, regardless of the choice of diagnostic or filters, the residual scatter is lower for asymptotically high values of α than it is for $\alpha = 0$. This implies that, in general, color is more effective than luminosity as a predictor of metallicity.

In contrast, the residual scatter achieved by Mannucci et al. (2010) in terms of the optimal projection of the physical parameters stellar mass (M_*) and SFR, $\mu_{0.32} = \log(M_*) - 0.32 \log(\text{SFR})$, was only 0.05 dex.




FIG. 2.— The optimal projection of the LZC relation for M_r , $g-r$ color, and three different metallicity diagnostics (T04, KD02, and PP04 O3N2). The projected LZC relation is shown for the optimal value of α , with the best fit third order polynomial LZC relation in black, extending over the calibrated range (2 – 98th percentile). The color coding shows the optical physical parameter ($\mu_{0.32}$) from Mannucci et al. (2010). The insets display the dispersion in the residuals in metallicity from the LZC as a function of the color-weighting parameter α , with the optimal value marked by the dashed line.

However, estimation of $\mu_{0.32}$ is based on full $ugriz$ imaging and $R \sim 2000$ optical spectroscopy, while the LZC relation relies on imaging in just two bands and a redshift estimate. The coloring in Figure 2 illustrates that the optimal projection of the photometric properties is highly correlated with $\mu_{0.32}$, with Pearson correlation coefficient $\rho \sim 0.8$ for all metallicity diagnostics and filters.

For two of the metallicity diagnostics we consider (P01 and PT05, both based on P), the strength of the LZC is negligible ($\sigma_Z/\sigma_{Z,0} \sim 1$). This could imply that the information expressed by galaxy color is already included in these metallicity diagnostics, such that they produce more physically realistic metallicity estimates than the other diagnostics. However, we find the opposite scenario to be more likely. These metallicity diagnostics have well known discrepancies with other strong line diagnostics, yielding the M-Z relation essentially unobservable (Kewley & Ellison 2008). Similarly, we find that they produce extreme scatter when used to calibrate the luminosity-metallicity relation ($\sigma_{Z,0} = 0.19$ dex for P01, M_r , $(r-i)$), which is not improved by the incorporation of color information. However, the LZC relation calibrated with the PVT diagnostic (also based on P) does exhibit significant decrease in scatter ($\sigma_Z/\sigma_{Z,0} \sim 0.85$ depending on filter set).

As expected, choosing a bluer filter set produces the greatest improvement in σ_Z because it better reflects the size of the young stellar population. Taking the median across all metallicity diagnostics and using M_i , we find $\sigma_Z/\sigma_{Z,0} = [0.81, 0.82, 0.88]$ for $[g-r, g-i, r-i]$. The choice of filter for L determination, among gri , has a less significant effect. Using the $g-i$ color, we find $\sigma_Z/\sigma_{Z,0} = [0.78, 0.80, 0.82]$ for $[g, r, i]$.

4. APPLICATIONS

The calibrated LZC relation we present could benefit several disciplines, allowing for precise and accurate metallicity estimates for galaxies based on photometry alone. Spectroscopic metallicity measurement demands considerably more observational resources, while full SED modeling (e.g. Pacifici et al. 2012) is computationally demanding. In contrast, the LZC relation can be employed using existing multi-band photometry from wide-field surveys such as SDSS or newly acquired, targeted observations, so long as an estimate for the redshift of each galaxy is available. Using $ugriz$ photometry, photometric redshift estimates for galaxies can be achieved with scatter $\delta z \sim 0.02$ (Ball et al. 2008), while K -corrections can be determined to $\lesssim 20\%$ (Blanton & Roweis 2007).

To test the uncertainty in metallicity introduced by use of photometric redshifts (photo- z), we recompute metallicities for the subset of $\sim 70,000$ galaxies in the MPA-JHU catalog with photo- z estimates in the SDSS-DR8. We use the kd -tree nearest neighbor fit photo- z estimates, as described in Csabai et al. (2007). We compute K -corrections using the analytic prescriptions of Chilingarian et al. (2010), using both the spectroscopic and photometric redshifts for each galaxy. We apply the LZC relation as calibrated for the PP04 O3N2 metallicity diagnostic using the M_g luminosities and $g-r$ colors. The resulting distribution of metallicity residuals (δZ) for the spectroscopic versus photometric redshifts suggests that there is no systematic bias introduced by photo- z (median $\delta Z = 0.000$ dex). The typical uncertainty added to the metallicity estimates is negligible (standard deviation $\delta Z = 0.008$ dex) and therefore the photometric metallicity estimate is dominated by the

TABLE 2
PARAMETERS OF THE *LZC* RELATIONS

<i>L</i>	Color	α	$p_3 \times 10^5$	$p_2 \times 10^3$	$p_1 \times 10^3$	p_0	σ_Z	$\sigma_{Z,0}$	$\sigma_Z/\sigma_{Z,0}$	μ range ^a
PP04 O3N2										
<i>g</i>	<i>g</i> − <i>r</i>	28.7	0.80	−1.02	−122.93	6.01	0.130	0.088	0.678	[−43.3, −26.5]
<i>g</i>	<i>g</i> − <i>i</i>	11.8	10.00	5.13	16.54	7.33	0.130	0.088	0.677	[−34.0, −22.4]
<i>g</i>	<i>r</i> − <i>i</i>	18.1	2.88	0.81	−53.50	7.37	0.130	0.099	0.761	[−26.9, −19.1]
<i>r</i>	<i>g</i> − <i>r</i>	27.7	0.80	−1.02	−122.93	6.01	0.127	0.088	0.696	[−43.3, −26.5]
<i>r</i>	<i>g</i> − <i>i</i>	10.9	11.97	6.42	42.01	7.47	0.127	0.088	0.691	[−33.7, −22.4]
<i>r</i>	<i>r</i> − <i>i</i>	15.9	4.15	1.10	−63.26	7.14	0.127	0.098	0.773	[−26.8, −19.4]
<i>i</i>	<i>g</i> − <i>r</i>	24.2	2.47	0.14	−97.70	6.21	0.123	0.088	0.712	[−40.8, −25.6]
<i>i</i>	<i>g</i> − <i>i</i>	10.8	10.00	5.13	16.56	7.33	0.123	0.088	0.714	[−34.0, −22.4]
<i>i</i>	<i>r</i> − <i>i</i>	14.9	4.15	1.10	−63.26	7.14	0.123	0.098	0.795	[−26.8, −19.4]

NOTE. — Parameters of the *LZC* relation defined in Equation 1; *L* is the photometric band of the luminosity, α is the optimal value of the color weighting factor to minimize the scatter in metallicity, and p_i are the parameters of the best-fit *LZC* polynomial for the corresponding value of α . The value $\sigma_Z/\sigma_{Z,0}$ expresses the reduction in the scatter relative to the *LZ* relation (without color term). Table 2 is published in its entirety in the electronic edition and at <https://www.cfa.harvard.edu/~nsanders/papers/LZC/summary.htm>, including parameters for all metallicity diagnostics. A portion is shown here for guidance regarding its form and content.

^a The range of μ , the optimal projection of luminosity and color, over which the diagnostic is calibrated. This range is defined by the 2nd and 98th percentiles in μ of the SDSS galaxies in the calibration sample.

scatter in the *LZC* relation.

Lara-López et al. (2010) concluded that there is no detectable evolution in the fundamental plane for star-forming galaxies out to $z \sim 3.5$. Similarly, Cresci et al. (2012) inspect ~ 300 zCOSMOS galaxies and find no evolution in the Fundamental Metallicity Relation to $z \sim 0.8$. This would imply that observers who wish to apply the *LZC* relation to imaging of galaxies in this redshift regime need only account for *K*-corrections. However, Perez-Montero et al. (2012) investigate a larger sample of ~ 5000 zCOSMOS galaxies to $z \sim 1.3$ and report evolution of the SFR-corrected mass-metallicity relation starting at $z \gtrsim 0.4$. This suggests that additional studies are needed to calibrate the *LZC* for redshifts $z \gtrsim 0.4$.

Studies of supernova (SN) host galaxies can support inferences into progenitor star populations (Modjaz et al. 2008; Prieto et al. 2008; Sanders et al. 2012b, e.g.), with some studies relying on photometry rather than spectroscopy to measure the host galaxy metallicity (Prantzos & Boissier 2003; Boissier & Prantzos 2009; Arcavi et al. 2010, e.g.). However, Sanders et al. (2012b) have shown that the statistical uncertainty associated with metallicity estimates based on the galaxy luminosity-metallicity relation is a significant barrier to detecting subtle differences in metallicity among SN populations. Moreover, because some SNe strongly prefer blue galaxies (high SFR environments; Levesque et al. 2010; Kelly & Kirshner 2011), photometric metallicity estimates will be biased if color information is not incorporated. Using the *LZC* relation will effectively remove this bias, and significantly reduce the uncertainty in metallicity measurements. The additional uncertainty introduced by photo-*z* should be minor, as SN host galaxies studies are almost exclusively done in the $z < 0.15$ regime (see compilation in Sanders et al. 2012b), and spectroscopic redshift estimates are often available from the SN spectroscopy. We note that the *LZC* relation predicts the “global” or nuclear galaxy metallicity, as probed by the 3” SDSS spectroscopic fibers, and significant offsets may exist from the SN host environment metallic-

ity due to metallicity gradients in galaxies. However, these metallicity offsets are typically small ($\lesssim 0.1$ dex; Sanders et al. 2012b), and the intrinsic scatter in galaxy metallicity profiles limits the ability to isolate the explosion site even with spectroscopy (Rosolowsky & Simon 2008; Sanders et al. 2012a).

Similarly, galaxy metallicity measurements are key to the discussion of the progenitor properties of long-duration gamma-ray bursts (LGRBs, see e.g. Fynbo et al. 2003; Sanders et al. 2012c). While the host environments of LGRBs typically fall below the mass-metallicity relation (Levesque et al. 2010), it has been shown that these host galaxies do follow the fundamental plane relation (Mannucci et al. 2011, but see also Kocevski & West 2011). Because LGRBs are frequently discovered at high redshift ($z > 1$), the *LZC* relation could be used to derive metallicity estimates at considerably lower expense than deep NIR spectroscopy (e.g. Maiolino et al. 2008). However, in order to study the extremely lowest metallicity environments preferred by LGRBs ($Z \lesssim 0.3 Z_\odot$), additional calibration is needed to extend the *LZC* relation beyond the range probed by the SDSS galaxy sample, $0.4 Z_\odot \lesssim Z \lesssim 1.3 Z_\odot$ (see e.g. Mannucci et al. 2011).

As a usage example, we apply the *LZC* relation to the unusual host galaxy of the SN 2010ay. In Sanders et al. (2012c), we report that this host galaxy is a 2σ outlier from the luminosity-metallicity relation. The median and 1σ metallicity interval for SDSS galaxies with luminosity similar to this host galaxy ($M_B = -18.3$ mag) is $12 + \log(\text{O}/\text{H}) = 8.93 \pm 0.17$ (T04). This is a factor of $\gtrsim 2$ greater than the spectroscopically-measured T04 metallicity of $12 + \log(\text{O}/\text{H}) = 8.58$. The discrepancy is due to the extremely low mass-to-light ratio and high SFR of the host galaxy. Using the *LZC* relation for M_r and $r - i$ color, we find a T04 metallicity of $12 + \log(\text{O}/\text{H}) = 8.72 \pm 0.09$. This agrees with the spectroscopically measured value to within $\sim 40\%$ and has a significantly lower associated uncertainty. However, the *LZC* relation cannot be applied in all filter combinations because the host lies outside the calibrated range for μ ,

illustrating the importance of extending the calibration presented here to lower-metallicity host galaxies not well-represented in the SDSS spectroscopic sample.

Finally, we suggest that the next generation of wide field, multi-band, photometric surveys could use the *LZC* relation to characterize the metallicity distribution of galaxies in the local universe, and its evolution with redshift. The Pan-STARRS1 (PS1) survey is already operating, and will provide g_{P1} r_{P1} i_{P1} z_{P1} y_{P1} photometry for $\sim 2 \times 10^8$ galaxies over 3/4 of the sky (Saglia et al. 2012). Simultaneously, the Baryon Oscillation Spectroscopic Survey (BOSS Dawson et al. 2012) is providing spectroscopic redshifts to $\sim 1.5 \times 10^6$ galaxies to $z \sim 0.7$, but with emission line S/N typically too low for spectroscopic metallicity analysis. In the future, the Large Synoptic Survey Telescope (LSST) will provide *ugrizy* photometry of $\sim 10^{10}$ galaxies to $z \sim 6$ (LSST Science Collaboration et al. 2009). With the advent of such datasets, the *LZC* relation may play an

important role in defining the metallicity distribution of galaxies that has emerged from the cosmic evolution of star formation and galaxy mass.

We thank the MPA/JHU teams for making available their catalog of measured properties for SDSS galaxies, and we thank D. Eisenstein, D. Erb, R. Foley, and A. Tripathi for helpful conversations. This work was supported by the National Science Foundation through a Graduate Research Fellowship provided to N.E.S.; E.M.L. is supported by NASA through Einstein Postdoctoral Fellowship grant number PF0-110075 awarded by the Chandra X-ray Center, which is operated by the Smithsonian Astrophysical Observatory for NASA under contract NAS8-03060; and support for this work was provided by the David and Lucile Packard Foundation Fellowship for Science and Engineering awarded to A.M.S.

Facilities: PS1, EVLA, Swift, MMT

REFERENCES

- Abazajian, K. N., et al. 2009, *ApJS*, 182, 543
 Arcavi, I., et al. 2010, *ApJ*, 721, 777
 Ball, N. M., Brunner, R. J., Myers, A. D., Strand, N. E., Alberts, S. L., & Tcheng, D. 2008, *ApJ*, 683, 12
 Bell, E. F., & de Jong, R. S. 2001, *ApJ*, 550, 212
 Blanton, M. R., & Roweis, S. 2007, *AJ*, 133, 734
 Boissier, S., & Prantzos, N. 2009, *A&A*, 503, 137
 Brinchmann, J., Charlot, S., White, S. D. M., Tremonti, C., Kauffmann, G., Heckman, T., & Brinkmann, J. 2004, *MNRAS*, 351, 1151
 Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245
 Chilingarian, I. V., Melchior, A.-L., & Zolotukhin, I. Y. 2010, *MNRAS*, 405, 1409
 Cresci, G., Mannucci, F., Sommariva, V., Maiolino, R., Marconi, A., & Brusa, M. 2012, *MNRAS*, 421, 262
 Csabai, I., Dobos, L., Trencsényi, M., Herczegh, G., Józsa, P., Purger, N., Budavári, T., & Szalay, A. S. 2007, *Astronomische Nachrichten*, 328, 852
 Dawson, K. S., et al. 2012, *ArXiv e-prints*, 1208.0022
 Fynbo, J. P. U., et al. 2003, *A&A*, 406, L63
 Garnett, D. R., & Shields, G. A. 1987, *ApJ*, 317, 82
 Kauffmann, G., et al. 2003a, *MNRAS*, 346, 1055
 ——. 2003b, *MNRAS*, 341, 33
 Kelly, P. L., & Kirshner, R. P. 2011, *ArXiv e-prints*, 1110.1377
 Kewley, L. J., & Dopita, M. A. 2002, *ApJS*, 142, 35
 Kewley, L. J., & Ellison, S. L. 2008, *ApJ*, 681, 1183
 Kobulnicky, H. A., Kennicutt, Jr., R. C., & Pizagno, J. L. 1999, *ApJ*, 514, 544
 Kobulnicky, H. A., & Kewley, L. J. 2004, *ApJ*, 617, 240
 Kocevski, D., & West, A. A. 2011, *ApJ*, 735, L8+
 Lara-López, M. A., et al. 2010, *A&A*, 521, L53+
 Lequeux, J., Peimbert, M., Rayo, J. F., Serrano, A., & Torres-Peimbert, S. 1979, *A&A*, 80, 155
 Levesque, E. M., Kewley, L. J., Berger, E., & Zahid, H. J. 2010, *AJ*, 140, 1557
 Lopez-Sanchez, A. R., Dopita, M. A., Kewley, L. J., Zahid, H. J., Nicholls, D. C., & Scharwachter, J. 2012, *ArXiv e-prints*, 1203.5021
 LSST Science Collaboration, et al. 2009, *ArXiv e-prints*, 0912.0201
 Maiolino, R., et al. 2008, *A&A*, 488, 463
 Mannucci, F., Cresci, G., Maiolino, R., Marconi, A., & Gnerucci, A. 2010, *MNRAS*, 408, 2115
 Mannucci, F., Salvaterra, R., & Campisi, M. A. 2011, *MNRAS*, 439
 McGaugh, S. S. 1991, *ApJ*, 380, 140
 Modjaz, M., et al. 2008, *AJ*, 135, 1136
 Osterbrock, D. E., & Ferland, G. J. 2006, *Astrophysics of gaseous nebulae and active galactic nuclei*, ed. Osterbrock, D. E. & Ferland, G. J.
 Pacifici, C., Charlot, S., Blaizot, J., & Brinchmann, J. 2012, *MNRAS*, 421, 2002
 Peeples, M. S., & Shankar, F. 2011, *MNRAS*, 417, 2962
 Perez-Montero, E., et al. 2012, *ArXiv e-prints*, 1210.0334
 Pettini, M., & Pagel, B. E. J. 2004, *MNRAS*, 348, L59
 Pilyugin, L. S. 2001, *A&A*, 374, 412
 Pilyugin, L. S., & Thuan, T. X. 2005, *ApJ*, 631, 231
 Pilyugin, L. S., Vilchez, J. M., & Thuan, T. X. 2010, *ApJ*, 720, 1738
 Prantzos, N., & Boissier, S. 2003, *A&A*, 406, 259
 Prieto, J. L., Stanek, K. Z., & Beacom, J. F. 2008, *ApJ*, 673, 999
 Rosolowsky, E., & Simon, J. D. 2008, *ApJ*, 675, 1213
 Saglia, R. P., et al. 2012, *ApJ*, 746, 128
 Salim, S., et al. 2007, *ApJS*, 173, 267
 Sanders, N. E., Caldwell, N., McDowell, J. C., & Harding, P. 2012a, *ArXiv e-prints*, 1209.2251
 Sanders, N. E., et al. 2012b, *ArXiv e-prints*, 1206.2643
 ——. 2012c, *ApJ*, 756, 184
 Tremonti, C. A., et al. 2004, *ApJ*, 613, 898